

GONORRHEA and CHLAMYDIA

HEALTH CARE PROVIDER INVESTIGATION & REPORTING FORM

The Health Protection and Promotion Act 1990 (HPPA), R.S.O., 1990, and Ontario Reg. 135/18, outlines the requirements for physicians, practitioners, and institutions to report any **disease of public health significance** to the Medical Officer of Health.

Completion of both pages of this form is required. The form is to be faxed by the next working day from the initial patient visit, to the Windsor-Essex County Health Unit (WECHU) – Infectious Disease Prevention Department (fax: 226-783-2132). Refer to the *Canadian Guidelines on Sexually Transmitted Infections (STIs)* for diagnosis and management of STIs.

DATE REPORTED (YY/MM/DD)		REPORTING PROVIDER NAME		PHONE NUMBER		ext.		
CECTION A	- DATIENT INCORNATION				() -		ext.	
SECTION A: PATIENT INFORMATION								
PATIENT NAME				SEX	DATE OF BIRTH (YY/MM	M/DD)	AGE	
	(FIRST) (MIDDLE)	(LAST)					
ADDRESS								
(STREET)			(CITY) (POSTAL CODE)					
HOME PHONE: () -				ALTERNATE PHO	IE: (-			
☐ Yes ☐ No	Has the client been notified of the laboratory result, indicating infection?							
☐ Yes ☐ No	Is the client pregnant? If yes, indicate gestational age: weeks							
SECTION B: PRESENTING SYMPTOMS								
✓ FEMALES			Onset Date YY/MM/DD)	✓ MALES	ES		Onset Date (YY/MM/DD)	
☐ Asymptomatic (most common)				☐ Asymptomatic				
☐ Lower abdominal pain				☐ Dysuria				
☐ Deep dyspareunia				☐ Testicular pain	ain			
☐ Dysuria				☐ Urethral discha	l discharge			
☐ Rectal pain/discharge and proctitis			☐ Urethral itch					
☐ Abnormal vaginal bleeding			☐ Rectal pain/dis	scharge and proctitis				
☐ Vaginal discharge			☐ Other, specify:					
☐ Other, specify:								
SECTION C: RISKS FOR INFECTION AND COMPLICATIONS								
✓ RISKS								
\square Sexual contact of a confirmed gonorrhea or chlamydia case \square Multiple sex partners								
☐ Sex with same sex			☐ New sexual contact in the past 2 months					
☐ Sex with opposite sex			☐ Alcohol and/or drug use					
□ No condom use			☐ Those with street involvement/homeless					
☐ Condom breakage			☐ Unprotected sea	☐ Unprotected sex while traveling to endemic area				
☐ Anonymous sex partners			☐ Sex trade worker					

SECTION D	: INFECTION MANAGEMENT							
☐ Yes	Was treatment provided to the client? If yes, specify medication & date below.							
□ No	All confirmed cases need to be treated and suspected cases should be considered for treatment.							
	NOTE : Free STIs medications can be ordered from the WECHU to have in your office for prompt treatment.							
	** NEW - TREATMENT PER GUIDELINES FOR UNCOMPLICATED GONORRHEA INFECTIONS AND CHLAMYDIA IN ADULTS							
	First-line Treatment ☐ Ceftriaxone 500mg IM single dose	DATE GIVEN (YY/MM/DD):						
	AND							
	☐ Azithromycin 1 g PO single dose							
	Alternative Treatments	DATE GIVEN (YY/MM/DD):						
	☐ Cefixime 800mg PO AND *Doxycycline 100mg PO BID x 7 days							
	(*Not for use in pregnant/lactating individuals) OR							
	☐ Cefixime 800mg PO AND Azithromycin 1g PO							
	☐ Other:	DATE GIVEN (YY/MM/DD):						
	a other.							
☐ Yes ☐ No	Advise client to inform sexual partners to see a health care provider for testing and treatment. Inform client that WECHU can assist with anonymous partner notification.							
	WEETTO can assist with anonymous partier notineation.							
#:	# of sexual partners identified by the client 60 days prior.							
SECTION E	: CLIENT EDUCATION							
	Counsel client regarding transmission and prevention methods. Advise client/contact(s) to abstain from or have protected intercourse of all types (anal, oral, and vaginal) until at least 7 days after completion of appropriate treatment and the clients/contact(s) are asymptomatic.							
	** NEW - Test of Cure (TOC) is recommended for all positive Gonorrhea sites in all cases. This is particularly important when regimens other than ceftriaxone 500mg IM are used.							
	Obtain NAAT three to four weeks after completion of treatment OR obtain culture at least three days after completion of treatment. If treatment failure is suspected more than three weeks after treatment (e.g., when symptoms persist or recur after treatment), complete both NAAT and Culture.							
	Inform client that repeat testing for gonorrhea is recommended 6 months post-treatment, as reinfection is high.							
	Inform client that a nurse from the WECHU may be contacting them.							
* The Public Health Lab Service Desk (1-877-604-4567) is available to answer questions regarding specimen collection. An online test information index is also available at www.publichealthontario.ca .								
REPORTIN	REPORTING HEALTH CARE PROVIDER'S SIGNATURE:							

Client Name: _____ Client DOB:_____

The most current form is available on our website:

https://www.wechu.org/forms/

GC/Chlamydia Page 2 of 2